arXiv:1611.05423v1 [math.CO] 16 Nov 2016

Density of monochromatic infinite subgraphs

Louis DeBiasio[1], Paul McKenney[2]

November 17, 2016

Abstract

For any countably infinite graph G, Ramsey's theorem guarantees an infinite monochromatic copy of G in any r-coloring of the edges of the countably infinite complete graph $K_{\mathbb{N}}$. Taking this a step further, it is natural to wonder how "large" of a monochromatic copy of G we can find with respect to some measure – for instance, the density (or upper density) of the vertex set of G in the positive integers. Unlike finite Ramsey theory, where this question has been studied extensively, the analogous problem for infinite graphs has been mostly overlooked.

In one of the few results in the area, Erdős and Galvin proved that in every 2-coloring of $K_{\mathbb{N}}$, there exists a monochromatic path whose vertex set has upper density at least $2/3$, but it is not possible to do better than $8/9$. They also showed that for some sequence $\epsilon_n \to 0$, there exists a monochromatic path P such that for infinitely many n, the set $\{1, 2, ..., n\}$ contains the first $(\frac{1}{3+\sqrt{3}} - \epsilon_n)n$ vertices of P, but it is not possible to do better than $2n/3$. We improve both results, in the former case achieving an upper density at least $3/4$ and in the latter case obtaining a tight bound of $2/3$. We also consider related problems for directed paths, trees (connected subgraphs), and a more general result which includes locally finite graphs for instance.

1 Introduction

Throughout the paper we let $K_{\mathbb{N}}$ be the complete graph on the positive integers and $\vec{K}_{\mathbb{N}}$ be the complete symmetric digraph on the positive integers. By r-coloring of G, we mean a partition of the edges of G into r parts (i.e. an r-edge-coloring). We say that a collection of subgraphs $\{H_1, H_2, \ldots, H_k\}$ covers (partitions) G if the collection of vertex sets $\{V(H_1), V(H_2), \ldots, V(H_k)\}$ forms a cover (partition) of the vertex set $V(G)$.

In 1928 Ramsey [27] proved (stated here only for graphs) that for any r-coloring of $K_{\mathbb{N}}$ there exists an infinite set $A \subseteq \mathbb{N}$ such that all of the edges with both endpoints in A have the same color. It is natural to wonder if Ramsey's theorem can be strengthened to say something more precise about the size of A; for instance, can we control the growth of the elements of A? It seems to be first noted by Erdős in 1964 [5] that no such result can hold. That is, there exist 2-colorings of $K_{\mathbb{N}}$ in which every red complete subgraph is finite and the gaps between consecutive vertices of any blue complete subgraph grow arbitrarily fast.

[1]Department of Mathematics, Miami University debiasld@miamioh.edu. Research supported in part by Simons Foundation Collaboration Grant # 283194.

[2]Department of Mathematics, Miami University mckennp2@miamioh.edu.

In the case of sparser graphs (such as paths), it may be possible to find monochromatic copies with controlled growth. In this paper, we prove a variety of such results surrounding the *upper density* of a set $A \subseteq \mathbb{N}$;

$$\bar{d}(A) = \limsup_{n\to\infty} \frac{|A \cap \{1, \ldots, n\}|}{n}.$$

Given a graph H with vertex set $V(H) \subseteq \mathbb{N}$, we define the *upper density* of H to be that of $V(H)$. The *lower density* of A, denoted $\underline{d}(A)$, is defined analogously in terms of the lim inf. We note that other more general notions of size, defined via *ideals*[1] over $\mathbb{N}$, have been studied in the context of Ramsey's theorem going back to the 1970's; the interested reader is directed to [17], [16], and [15].

We note that in order to meaningfully talk about the upper density of a monochromatic copy of an infinite graph G in an edge-colored countably-infinite complete graph K, we must fix a ordering of the vertices of K before the coloring of the edges of K is given; as otherwise, for every coloring of the edges of K, there exists an ordering of the vertices of K so that K has a complete monochromatic subgraph of density 1. So for the purposes of computing the upper density, we simply treat the vertex set of K as the positive integers with the natural ordering.

1.1 Paths

One of the first results in graph Ramsey theory is the following.

Theorem 1.1 (Gerencsér, Gyárfás [9])**.** *In every 2-coloring of K_n there exists a monochromatic path with more than $2n/3$ vertices. Furthermore, this is best possible .*

Towards an infinite analogue of Theorem 1.1, one might ask; given a 2-coloring of $K_{\mathbb{N}}$, what is the largest asymptotic density of a monochromatic path, i.e. a monochromatic copy of the graph P with ordered vertex set $V(P) = \langle x_n \mid n \in \mathbb{N}\rangle$ and edge set $E(P) = \{x_n x_{n+1} \mid n \in \mathbb{N}\}$?

Erdős and Galvin [7],[6] seem to be the first to consider this question. Among other things, they showed that there is a 2-coloring for which every monochromatic subgraph of bounded maximum degree (in particular, paths) has lower density 0 ([7, Theorem 1.1]). This means that we cannot expect results using the usual asymptotic density. On the other hand, a result of Rado [26] states that in any r-coloring of $K_{\mathbb{N}}$, there is a partition of $\mathbb{N}$ into at most r monochromatic paths, and this implies that one of these paths must have upper density at least $1/r$. In the case of 2 colors, Erdős and Galvin strengthened this lower bound, and also found an upper bound;

Theorem 1.2 (Erdős, Galvin [7])**.** *In every 2-coloring of $K_{\mathbb{N}}$ there exists a monochromatic path with upper density at least $2/3$. On the other hand, there exists a 2-coloring of $K_{\mathbb{N}}$ in which every monochromatic path has upper density at most $8/9$.*

Erdős and Galvin also considered an alternate notion of density of a path P which only measures the density of "unbroken" initial segments of P.

Definition 1.3. *Given an infinite sequence of distinct natural numbers $A = (a_1, a_2, \ldots, a_n, \ldots)$, the* strong upper density *of A and the* strong lower density *of A are defined to be*

$$\bar{d}_s(A) = \limsup_{n\to\infty} \frac{f(n)}{n} \quad \text{and} \quad \bar{d}_s(A) = \liminf_{n\to\infty} \frac{f(n)}{n}$$

respectively, where $f(n) = \sup\{m \mid \{a_1, \ldots, a_m\} \subseteq \{1, \ldots, n\}\}$.



Figure 1: A path P which has density $8/12$ in the set $\{1, 2, \ldots, 12\}$, but which has strong density $6/12$ in the set $\{1, 2, \ldots, 12\}$.

In other words, in the strong upper density of a sequence A we consider only the density of the longest initial segment of A which lies entirely inside each set $\{1, \ldots, n\}$ (see Figure 1). In particular, we have $\bar{d}_s(A) \leq \bar{d}(A)$. Given a graph G with an ordered vertex set $V(G) = (v_1, v_2, \ldots, v_n, \ldots)$ consisting of distinct natural numbers, we define the *strong upper density* of G to be the strong upper density of the sequence $(v_1, v_2, \ldots, v_n, \ldots)$.

Again for 2 colors, Erdős and Galvin offered the following bounds on the strong upper density of monochromatic paths;

Theorem 1.4 (Erdős, Galvin [7])**.** *In every 2-coloring of $K_{\mathbb{N}}$ there exists a monochromatic path with strong upper density at least $1/(3+\sqrt{3})$. On the other hand, there exists a 2-coloring of $K_{\mathbb{N}}$ in which every monochromatic path has strong upper density at most $2/3$.*

For $r \geq 3$ colors they proved a few results, including the somewhat surprising fact that there exists an r-coloring of $K_{\mathbb{N}}$ in which every monochromatic path has strong upper density 0 ([7, Theorem 1.7].

Theorem 1.5 (Erdős, Galvin [7])**.** *Let $r \geq 3$.*

(i) *There exists an r-coloring of $K_{\mathbb{N}}$ such that for all monochromatic paths P, $\bar{d}(P) \leq 2/r$.*

(ii) *There exists an r-coloring of $K_{\mathbb{N}}$ such that for all monochromatic paths P, $\bar{d}_s(P) = 0$.*

We improve the lower bound for ordinary upper density of paths.

Theorem 1.6. *In every 2-coloring of $K_{\mathbb{N}}$, there exists a monochromatic path P such that $\bar{d}(P) \geq 3/4$.*

Furthermore, we prove a tight result for the strong upper density of paths in 2-colored complete graphs. As this was the only open case, this completely settles the strong upper density problem for paths.

Theorem 1.7. *In every 2-coloring of $K_{\mathbb{N}}$, there exists a monochromatic path P such that $\bar{d}_s(P) \geq 2/3$.*

Finally, we provide an improved upper bound for the upper density of paths when $r \geq 3$ (Corollary 3.5).

1.2 Directed paths

In an orientation of a path $P = v_1 v_2 \ldots$, we say that v_i is an *out-(in-)switch* if the in-(out-)degree of v_i is 0 in P. Note that any endpoints of P are switches, and that P has one endpoint if P is infinite and two endpoints if P is finite. We say that P is consistently oriented if the only switches of P are the endpoints, and we say that P is anti-directed if every vertex in P is a switch.

Raynaud proved a directed analog of Theorem 1.1.

[1] An *ideal* over a set X is a collection $\mathscr{I}$ of subsets of X which is closed under taking subsets and closed under finite unions.

Theorem 1.8 (Raynaud [28]). *In every 2-coloring of $\vec{K}_n$, there exists a consistently oriented monochromatic path with at least $n/2+1$ vertices. Furthermore, this is best possible.*

As above with undirected graphs, it is natural to consider an infinite analog of Raynaud's theorem. So it is somewhat surprising that we prove that for all $\epsilon > 0$, there exists a 2-coloring of $\vec{K}_{\mathbb{N}}$ such that every consistently oriented monochromatic path has upper density less than ϵ (Example 3.1). And for $r \geq 3$, there exists an r-coloring of $\vec{K}_{\mathbb{N}}$ such that every monochromatic consistently oriented path has upper density 0 (Example 3.3).

As mentioned in the previous section, Rado proved that in every r-coloring of $K_{\mathbb{N}}$, there exists a partition of $\mathbb{N}$ into at most r monochromatic paths (which implies that some monochromatic path has upper density at least $1/r$). Actually, his result was more general. He proved that in every r-coloring of $\vec{K}_{\mathbb{N}}$, there exists a partition of $\mathbb{N}$ into at most r monochromatic anti-directed paths (which implies that some monochromatic anti-directed path has upper density at least $1/r$). So for digraphs, the density depends strongly on the orientation of the path.

We show that if P contains infinitely many switches, then in any r-coloring of the edges of $\vec{K}_{\mathbb{N}}$ (actually, in any r-coloring of the edges of any tournament on $\mathbb{N}$), there exists a monochromatic copy of P with positive upper density (Corollary 7.4). For paths with finitely many switches, the result is equivalent to the results above for consistently oriented paths.

1.3 Connected subgraphs

An old remark of Erdős and Rado says that a graph or its complement is connected, or alternatively, in every 2-coloring of K_n there exists a monochromatic connected subgraph on n vertices. This remark was extended by Gyárfás, and is one of the few examples in graph Ramsey theory for which a tight result is known for infinitely many values of r.

Theorem 1.9 (Gyárfás [10]). *Every r-coloring of K_n contains a monochromatic connected subgraph on at least $n/(r-1)$ vertices. Furthermore, this is best possible when $r-1$ is a prime power.*

We consider the analogous problem for infinite graphs. First we need a definition of strong upper density for connected graphs.

Definition 1.10. *Let T be a connected subgraph of $K_{\mathbb{N}}$. The* strong upper density *of T is*

$$\bar{d}_s(T) = \sup_{V \in \pi} \bar{d}_s(V)$$

where π is the set of all orderings $(v_1, v_2, \dots)$ of $V(T)$ such that for all $i \geq 1$, the graph $T[\{v_1, \dots, v_k\}]$ induced by $\{v_1, \dots, v_k\}$ is connected.

The reader can easily check that the set π in the above definition is nonempty. Moreover, we note that, for a connected subgraph T of $K_{\mathbb{N}}$, to show that $\bar{d}_s(T) = \alpha$ it suffices to find an increasing sequence of integers $n_1 < n_2 < \cdots < n_k < \cdots$ such that for each k, the subgraph of T induced by $V(T) \cap \{1, \dots, n_k\}$ is connected, and $\lim_{k\to\infty} |V(T) \cap \{1, \dots, n_k\}|/n_k = \alpha$.

It's not hard to see that the remark of Erdős and Rado easily extends to countably infinite graphs which shows that every 2-coloring of $K_{\mathbb{N}}$ contains a monochromatic connected subgraph T such that $V(T) = \mathbb{N}$, and thus has *density* 1; however, for $r \geq 3$ colors, Example 3.7 shows that upper density is all we can hope for. We show that the infinite analog of Erdős and Rado's remark extends to strong upper density, and Example 3.6 shows that we must consider strong upper density rather than ordinary strong density.

Theorem 1.11. *Every 2-coloring of $K_{\mathbb{N}}$ contains a monochromatic connected graph T such that $\bar{d}_s(T) = 1$.*

A (special case of a) conjecture of Ryser says that in every r-coloring of K_n, there exists a cover of $V(K_n)$ by at most $r-1$ monochromatic connected subgraphs, and if true this would imply Theorem 1.9. Ryser's conjecture is known to be true for $r \leq 5$ (see [30]). For $r = 3$, it is not hard to prove this result and see that it easily extends to countably infinite complete graphs. This implies that in every 3-coloring of $K_{\mathbb{N}}$ there exists a monochromatic connected subgraph with upper density at least $1/2$. It turns out to be considerably more difficult to show that the same holds for strong upper density, which we do for 3 colors.

Theorem 1.12. *Every 3-coloring of $K_{\mathbb{N}}$ contains a monochromatic connected graph T such that $\bar{d}_s(T) \geq 1/2$.*

As mentioned above, there exists an example to show that Theorem 1.9 is best possible when $r-1$ is a prime power. We show that this example extends to infinite graphs (Example 3.4).

1.4 0-dense graphs

For $r \geq 2$, we say that a graph G is $(0, r)$-*dense* if there exists an r-coloring of $K_{\mathbb{N}}$ such that every monochromatic copy of G has density 0, and if G is $(0, 2)$-dense, we just say that G is 0-*dense*. If for all $r \geq 2$, G is not $(0, r)$-dense, then we say that G is *positively-dense*. For directed graphs, we define $(0, r)$-dense and 0-dense in the analogous way. We show that if G has bounded independence number, then G is 0-dense (Proposition 7.1). We also show that if G has an infinite independent set B and every vertex in $V(G) \setminus B$ has finite degree to B, then G is positively-dense (Corollary 7.5).

2 Notation and outline

Given $n \in \mathbb{N}$ we will write $[n]$ for the set $\{1, \ldots, n\}$.

In many of the results to follow we will make use of a partition of $\mathbb{N}$ into consecutive intervals $A_1, A_2, \ldots, A_n, \ldots$ such that

$$\lim_{n\to\infty} \frac{|A_n|}{|A_1| + \cdots + |A_n|} = 1.$$

In such cases we will simply call the A_n's a sequence of *fast growing intervals*. In some situations we will require the sizes $|A_n|$ to grow even more rapidly, according to various conditions, in which case these conditions will be laid out explicitly.

Given a finite set $F \subseteq \mathbb{N}$, we define the *local density* of F to be $d_\ell(F) = |F|/\max(F)$. We note here that if A_n is a sequence of fast-growing intervals, then $d_\ell(A_n) \to 1$ as $n \to \infty$.

The paper is organized as follows. We collect all of our examples in Section 3. For the convenience of the reader we provide two of the examples of Erdős and Galvin, stated here in less general terms to suit our purpose, in the Appendix. In Section 4 we prove Theorem 1.6 which improves Erdős and Galvin's lower bound on the upper density of monochromatic paths in 2-colored $K_{\mathbb{N}}$. In Section 5 we prove the more complicated Theorem 1.7 which gives a tight bound on the strong upper density of monochromatic paths in 2-colored $K_{\mathbb{N}}$. This result uses the "regularity–blow-up method" based on Szemerédi's regularity lemma [29] and Komlós, Sárközy, and Szemerédi's blow-up lemma [22], and we collect all of the relevant background material at the beginning of that section. In Section 6 we prove Theorems 1.11 and 1.12 about the strong

upper density of monochromatic connected subgraphs in 2- and 3- colored $K_{\mathbb{N}}$. In Section 7 we discuss properties of graphs and digraphs which imply that they are 0-dense or positively dense. Finally, in Section 8, we discuss a large number of open questions stemming from our work.

3 Examples

3.1 Directed paths

Example 3.1. *For all $\epsilon > 0$, there exists a 2-coloring of $\vec{K}_{\mathbb{N}}$ such that every orientation of a monochromatic path with finitely many switches has upper density less than ϵ; in particular, this holds for consistently oriented paths.*

Proof. Let $\epsilon > 0$ and set $k = \lceil 1/\epsilon \rceil + 1$. Partition $\mathbb{N}$ into k infinite sets $A_1, \ldots, A_k$ each of density $1/k$ (for instance, partition based on residues modulo k). Let $x, y \in \mathbb{N}$. If $x, y \in A_i$ and $x < y$, then color (x, y) red and (y, x) blue. If $x \in A_i, y \in A_j$ and $i < j$, then color (x, y) blue and (y, x) red. Now let P be a monochromatic path with finitely-many switches. Then, P has infinite intersection with at most one of the sets A_i; thus P has upper density at most $1/k < \epsilon$. □

We note that the natural attempt to strengthen Example 3.1, by partitioning $\mathbb{N}$ into infinitely-many infinite sets $A_1, \ldots, A_i, \ldots$ each with density 0, fails due to the following fact.

Proposition 3.2. *Suppose $\mathbb{N} = A_1 \cup \cdots \cup A_i \cup \cdots$ is a partition of $\mathbb{N}$ into infinite sets, each with density 0. Then there is a set B with density 1 such that $B \cap A_i$ is finite for every i.*

Proof. Fix a sequence ϵ_i tending to zero. For each i, choose n_i large enough that for all $n \geq n_i$,

$$\frac{1}{n} \left| (A_1 \cup \cdots \cup A_i) \cap [n] \right| < \epsilon_i.$$

Now let

$$B = \bigcup_{i=1}^{\infty} A_i \cap [n_i]$$

Clearly each $B \cap A_i$ is finite; to see that B has density 1, note that for any given n, if $n_i \leq n < n_{i+1}$ then $[n] \setminus B \subseteq (A_1 \cup \cdots \cup A_i) \cap [n]$ and hence B has density at least $1 - \epsilon_i$ in $[n]$. □

In Problem 8.3 we ask whether one can find a 2-coloring of $\vec{K}_{\mathbb{N}}$ in which every monochromatic consistently-oriented path has density 0. We note here that 3 colors is enough to ensure not only that every such path has density 0, but that such paths grow arbitrarily fast.

Example 3.3. *Let $h : \mathbb{N} \to \mathbb{N}$, where $h(n) \to \infty$. Then there exists a 3-coloring of $\vec{K}_{\mathbb{N}}$ such that for every infinite monochromatic path $P = v_1 v_2 v_3 \ldots$ with finitely-many switches, the sequence $(v_1, v_2, \ldots)$ eventually grows faster than h.*

Proof. Let $\mathbb{N} = A_1 \cup A_2 \cup \cdots \cup A_n \cup \cdots$ be a partition of $\mathbb{N}$ into consecutive intervals, where $|A_n| \geq h(n)$. We color $\vec{K}_{\mathbb{N}}$ so that each A_n is complete in green, and for each $m < n$, the edges from A_m to A_n are red and the edges from A_n to A_m are blue. Let P be a monochromatic path with finitely-many switches. Then for all but finitely-many n, P intersects A_n in at most one vertex. It follows that for large enough k, the kth vertex of P is at least $h(k)$. □

3.2 Connected subgraphs

Example 3.4. *Let $r \geq 2$. If $r-1$ is a prime power, then there exists an r-coloring of $K_{\mathbb{N}}$ such that every monochromatic connected subgraph T has upper density at most $\frac{1}{r-1}$.*

Proof. Since $r-1$ is a power of a prime, there exists an affine plane of order $r-1$. Partition $\mathbb{N}$ into $(r-1)^2$ sets $V_0, \ldots, V_{(r-1)^2-1}$, each of density $1/(r-1)^2$ (for instance partition based on residues mod $(r-1)^2$). We will partition these $(r-1)^2$ sets into r parallel classes (one of each color) of $r-1$ "lines" each. Each line contains $r-1$ sets. Color the edges inside the sets arbitrarily.

So any monochromatic connected subgraph intersects at most $r-1$ of the sets $V_0, \ldots, V_{(r-1)^2-1}$ and thus has density at most $(r-1)/(r-1)^2 = 1/(r-1)$. □

Corollary 3.5. *Let $r \geq 3$ and let q be the largest prime power which is at most $r-1$. There exists an r-coloring of $K_{\mathbb{N}}$ such that every monochromatic path P has upper density at most $1/q < 2/r$.*

Proof. To verify that $1/q < 2/r$ for $r \geq 3$, first note that this holds for $r = 3$ and then let $r \geq 4$. By Bertrand's postulate, there exists a prime q' such that $\lfloor r/2 \rfloor < q' < 2\lfloor r/2 \rfloor$. Choosing q to be the largest such prime, we have $r/2 < q \leq r-1$ and thus $1/q < 2/r$. Now take the $(q+1)$-coloring of $K_{\mathbb{N}}$ given in Example 3.4, which is an r-coloring since $q+1 \leq r$. [2] □

The next two examples justify why we only consider (strong) upper density for monochromatic connected subgraphs.

Example 3.6. *There exists a 2-coloring of $K_{\mathbb{N}}$ such that every monochromatic connected subgraph T has strong lower density 0.*

Proof. Partition $\mathbb{N}$ into fast growing intervals $A_1, A_2, \ldots$. For all $x \in A_{2i+1}$ color all edges from x to $y \in A_j$ with $j \leq 2i+1$ red. For all $x \in A_{2i}$ color all edges from x to $y \in A_j$ with $j \leq 2i$ blue. If R is a red connected subgraph and B is a blue connected subgraph then

$$\underline{d}_s(R) = \lim_{i\to\infty} \frac{|A_1 \cup A_3 \cup \cdots \cup A_{2i-1}|}{|A_1 \cup A_2 \cup \cdots \cup A_{2i}|} = 0 \quad \text{and} \quad \underline{d}_s(B) = \lim_{i\to\infty} \frac{|A_2 \cup A_4 \cup \cdots \cup A_{2i}|}{|A_1 \cup A_2 \cdots \cup A_{2i+1}|} = 0$$

□

Example 3.7. *There exists a 3-coloring of $K_{\mathbb{N}}$ such that every monochromatic connected subgraph T has lower density 0.*

Proof. Partition $\mathbb{N}$ into fast growing intervals $A_0, A_1, A_2, \ldots$ and for each $q = 0, 1, 2$ and 3 let B_q be the union of all A_n such that $n \equiv q \pmod 4$. Note that each B_q has upper density 1. Let all edges between B_0 and B_3 and between B_1 and B_2 be red, all edges between B_0 and B_2 and between B_1 and B_3 be blue, and all edges between B_0 and B_1 and between B_2 and B_3 be green. Color all edges inside each B_q arbitrarily. Then any monochromatic, connected subgraph T is disjoint from some B_q; for instance any red connected subgraph must either be contained in $B_0 \cup B_3$ or $B_1 \cup B_2$. Hence any monochromatic connected subgraph has lower density 0. □

[2] Using stronger results than Bertrand's postulate, it is possible to get a better upper bound on $1/q$ for sufficiently large r.

4 Upper density of paths

We will use the following result implicitly proved by Gyárfás and Lehel (see Pokrovskiy [25, Corollary 1.9]).

Theorem 4.1 (Gyárfás, Lehel [11])**.** *In every 2-coloring of $K_{n,n}$ there exists a partition of the vertex set into at most three disjoint monochromatic paths.*

Let G be a graph in which the edges and vertices are colored with a set of r colors. A *monochromatic path forest* is a collection of vertex disjoint paths (i.e. a path forest) such that every edge and the endpoints of every path have the same color. Theorem 4.1 implies the following result about monochromatic path forests in 2-colored complete graphs.

Lemma 4.2. *Let a total 2-coloring of K_n be given where, if R is the set of red vertices and B the set of blue vertices, then $|R| \leq |B|$. Then for every $B' \subseteq B$ with $|B'| = |R|$, there exist red and blue monochromatic path forests F_R and F_B such that $F_R \subseteq R \cup B'$ and*

$$|F_R| + |F_B| \geq n + |R| - 3.$$

Proof. Let $B' \subseteq B$ with $|B'| = |R|$ and apply Theorem 4.1 to the bipartite graph induced by B' and R. If an endpoint of a blue path is in R or an endpoint of a red path is in B', then delete them to get blue paths with both endpoints in B' and red paths with both endpoints in R, and note that at most 3 endpoints are removed from B'. Letting F_R be the vertices in R together with the red paths and letting F_B be the vertices in B together with the blue paths gives

$$|F_R| + |F_B| \geq |B| + |R| + |B'| - 3 = n + |R| - 3,$$

as desired. □

Now we will use the Lemma above with a "density increment" argument to get a monochromatic path forest with density arbitrarily close to 3/4.

Lemma 4.3. *Let $0 < \epsilon \leq 3/4$ and $k \in \mathbb{N}$ with $k \geq 3/\epsilon$. For all $n \geq 4k/\epsilon$ and for every total 2-coloring of the complete graph K_n on the vertex set $[n]$, there exists a monochromatic path forest F such that for some $\ell \geq k$, $|F \cap [\ell]| \geq (3/4 - \epsilon)\ell$.*

Proof. Fix a total 2-coloring of the complete graph on the vertex set $[n]$, and let R and B be the sets of red vertices and blue vertices respectively. We will assume without loss of generality that $|R| \leq |B|$.

We may assume that for each $\ell \geq k$, $|R \cap [\ell]|$ is less than $|B \cap [\ell]|$; otherwise there is some $\ell \geq k$ such that $|R \cap [\ell]| = |B \cap [\ell]|$, and then by Lemma 4.2 there is a monochromatic path forest $F \subseteq [\ell]$ such that

$$|F| \geq \frac{1}{2}\left(\ell + \frac{\ell}{2} - 3\right) = \left(\frac{3}{4} - \frac{3}{2\ell}\right)\ell \geq \left(\frac{3}{4} - \epsilon\right)\ell.$$

Let $r_0 = |R|$ and for each i such that $r_i > 0$, let $b_{i+1} = r_i$ and let r_{i+1} be the number of red vertices before the b_{i+1}st blue vertex, according to the natural order of $[n]$. For each $i \geq 0$, let $J_i = [r_i + b_i]$. Note that J_i has exactly r_i red vertices and exactly b_i blue vertices; hence, by the assumption made above, for each i we have $r_{i+1} < b_{i+1} = r_i$. It follows that the sizes of the intervals J_i are decreasing. Let t be the largest integer such that $|J_t| \geq k$. Note that by the maximality of t, we must have $r_t < k$, as otherwise $b_{t+1} = r_t \geq k$ and thus $r_{t+1} + b_{t+1} \geq k$.

Moreover, if $b_t \geq 3r_t$, then the blue vertices in J_t form a monochromatic path forest with density at least $3/4$ in J_t; hence we may assume that

$$b_t - r_t < 2r_t < 2k. \tag{1}$$

The purpose of the following claim is to show that, if there is no monochromatic path forest with the desired density in some interval J_i, then the differences $b_i - r_i$ decrease as i increases. Combined with (1), it will follow that $b_0 - r_0$ is very small relative to n, and we will get a sufficiently dense monochromatic path forest by applying Lemma 4.2 again.

Claim 4.4. *For each i with $1 \leq i \leq t$, either*

(i) *there is a blue monochromatic path forest F_B with $|F_B \cap J_{i-1}| \geq (3/4 - \epsilon)|J_{i-1}|$,*

(ii) *there is a red monochromatic path forest F_R with $|F_R \cap J_i| \geq (3/4 - \epsilon)|J_i|$, or*

(iii) $b_{i-1} - r_{i-1} < b_i - r_i$.

Proof. Let $1 \leq i \leq t$ and consider the interval J_{i-1}. By Lemma 4.2, there are red and blue monochromatic path forests F_R and F_B contained in J_{i-1} such that $|F_R| + |F_B| = b_{i-1} + 2r_{i-1} - 3$, and $F_R \cap B$ is contained in the first $b_i = r_{i-1}$-many blue vertices of J_{i-1}. If $|F_B| \geq (3/4 - \epsilon)|J_{i-1}|$ then we are done. Assuming otherwise, we have

$$\begin{aligned} |F_R \cap B| = b_{i-1} + 2r_{i-1} - |F_B| - |F_R \cap R| - 3 &\geq b_{i-1} + r_{i-1} - |F_B| - 3 \\ &> \left(\frac{1}{4} + \epsilon\right)(b_{i-1} + r_{i-1}) - 3 \\ &\geq \frac{1}{4}(b_{i-1} + r_{i-1}). \end{aligned}$$

Now consider the size of $F_R \cap J_i$. Since $F_R \cap B \subseteq J_i$, we have

$$|F_R \cap J| > r_i + \frac{1}{4}(b_{i-1} + r_{i-1}).$$

If $|F_R \cap J_i| \geq (3/4 - \epsilon)|J_i|$, then we are done. Assuming otherwise, then, we get

$$r_i + \frac{1}{4}(b_{i-1} + r_{i-1}) < \frac{3}{4}(b_i + r_i)$$

Rearranging, we get the desired inequality $b_{i-1} - r_{i-1} < b_i - r_i$. □

By Claim 4.4, we may assume that $b_{i-1} - r_{i-1} < b_i - r_i$ for each i with $1 \leq i \leq t$, and thus by (1), we have $b_0 - r_0 < b_t - r_t < 2k$. So by Lemma 4.2, we may find a monochromatic path forest $F \subseteq J_0 = [n]$ with size at least

$$\frac{1}{2}(|B| + |R| + |R| - 3) \geq \frac{1}{2}(3|B| - 4k - 3) \geq \frac{1}{2}\left(\frac{3n}{2} - \epsilon n - 3\right) > \left(\frac{3}{4} - \epsilon\right)n.$$

This completes the proof. □

Proof of Theorem 1.6. Fix a 2-coloring of the edges of $K_{\mathbb{N}}$. We define a 2-coloring of the vertices by coloring $n \in \mathbb{N}$ red if and only if there are infinitely-many $m \in \mathbb{N}$ such that the edge $\{n, m\}$ is red.

Case 1:[3] there are vertices x and y of the same color, say red, and a finite set $S \subseteq \mathbb{N}$ such that there is no red path disjoint from S which connects x to y. We partition $\mathbb{N} \setminus S$ into sets X, Y, Z, where $x' \in X$ if and only if there is a red path, disjoint from S, which connects x to x', and $y' \in Y$ if and only if there is a red path disjoint from S which connects y to y'. Note that every edge from $X \cup Y$ to Z is blue. Since x and y are colored red, both X and Y are infinite, and by choice of x and y all edges in the bipartite graph between X and $Y \cup Z$ are blue. Hence there is a blue path with vertex set $X \cup Y \cup Z = \mathbb{N} \setminus S$.

Case 2: for every pair of vertices x and y of the same color χ, and every finite set $S \subseteq \mathbb{N}$, there is a path from x to y of color χ which is disjoint from S. Let ϵ_n be a sequence of positive reals tending to zero, and let a_n and k_n ($n \in \mathbb{N}$) be increasing sequences of integers such that

- $k_n \geq 3/\epsilon_n$ and $a_n \geq 4k_n/\epsilon_n$, and
- as $n \to \infty$, $k_n/(a_1 + \cdots + a_{n-1} + k_n) \to 1$.

Let $\mathbb{N} = A_1 \cup A_2 \cup \cdots \cup A_n \cup \cdots$ be a partition of $\mathbb{N}$ into consecutive intervals such that $|A_n| = a_n$. By Lemma 4.3 we may find monochromatic path forests $F_n \subseteq A_n$ and initial segments $I_n \subseteq A_n$ of length $\geq k_n$ such that

$$|F_n \cap I_n| \geq \left(\frac{3}{4} - \epsilon_n\right) |I_n| .$$

It follows that for any $B \subseteq \mathbb{N}$ containing infinitely-many F_n's, we have

$$\bar{d}(B) \geq \limsup_{n\to\infty} \frac{|F_n \cap I_n|}{a_1 + \cdots + a_{n-1} + |I_n|} = \limsup_{n\to\infty} \frac{3}{4} - \epsilon_n = \frac{3}{4}$$

By the pigeonhole principle, there are infinitely-many F_n's of the same color, say blue. We will recursively construct a blue path P which contains infinitely-many of these F_n's. To see how this is done, suppose we've constructed a finite initial segment p of P. We will assume as an inductive hypothesis that p ends on one of the end-vertices of some blue F_m. Let v be this end-vertex. Let n be large enough that $\min A_n$ is greater than every vertex in p, and F_n is blue. Let $w_1, \ldots, w_s$ list the end-vertices of F_n. By the case assumption, there is a blue path q_1 connecting v to w_1, such that q_1 is disjoint from $A_1 \cup \cdots \cup A_n$. Similarly, there is a blue path q_2 connecting w_1 to w_2, such that q_2 is disjoint from $A_1 \cup \cdots \cup A_n \cup q_1$. Continue in this fashion, choosing disjoint blue paths $q_3, \ldots, q_s$ such that q_i connects w_{i-1} to w_i. Clearly the paths of F_n allow us to put all of these together into a blue path p' which extends p and contains all of the vertices of F_n. □

Remark 4.5. *One may use an ultrafilter in the proof of Theorem 1.6 to define the 2-coloring of $\mathbb{N}$ in such a way that Case 1 never occurs. (See Theorem 7.3 for an example of this type of argument in action.) While this shortens the proof somewhat, the above proof has the advantage of being more elementary.*

5 Strong upper density of paths

To prepare for the proof of the main result in Section 5.3, we first develop a number of tools in the next two subsections.

5.1 Regularity and general lemmas

We will use the following version of Szemerédi's regularity lemma. For the basic definitions we refer the reader to [20] or [23].

[3] The proof in this case is exactly as in [7, Theorem 3.5].

Lemma 5.1 (2-colored regularity lemma – degree form)**.** *Let G be a 2-colored graph on n vertices. For all $0 < \epsilon \leq 1$ and positive integers m, there exists an $M = M(\epsilon, m)$ such that if $0 \leq d \leq 1$, then there is $m \leq k \leq M$, a partition $\{V_0, V_1, \ldots, V_k\}$ of the vertex set V and a subgraph $G' \subseteq G$ with the following properties:*

(i) $|V_0| \leq \epsilon n$,

(ii) *all clusters $V_1, \ldots, V_k$ are of the same size $|V_1| \leq \lceil \epsilon n \rceil$,*

(iii) $d_{G'}(v) > d_G(v) - (2d + \epsilon)n$ *for all $v \in V$,*

(iv) $e(G'[V_i]) = 0$ *for all $i \in [k]$,*

(v) *for all $1 \leq i < j \leq k$, the pair (V_i, V_j) is ϵ-regular in G'_1 with a density either 0 or greater than d and ϵ-regular in G'_2 with a density either 0 or greater than d, where $E(G') = E(G'_1) \cup E(G'_2)$ is the induced 2-coloring of G'.*

Definition 5.2 ((ϵ, d)-reduced graph)**.** *Given a graph G and a partition $\{V_0, V_1, \ldots, V_k\}$ satisfying conditions (i)-(v) of Lemma 5.1, we define the (ϵ, d)-reduced graph of G to be the graph Γ on vertex set $\{V_1, \ldots, V_k\}$ such that V_iV_j is an edge of Γ if $G'[V_i, V_j]$ has density greater than $2d$. For each $V_iV_j \in E(\Gamma)$, we assign color 1 if $G'_1[V_i, V_j]$ has density greater than d and color 2 if $G'_2[V_i, V_j]$ has density greater than d (note that since the total density is greater than $2d$ every edge must receive a color, but it need not be unique).*

The following two Propositions are commonly used together with the regularity lemma (see [20, Proposition 42 and 43]).

Proposition 5.3. *Let $0 < 2\epsilon \leq d \leq c/2$ and let G be a graph on n vertices with $\delta(G) \geq cn$. If Γ is a (ϵ, d)-reduced graph of G obtained by applying Lemma 5.1, then $\delta(\Gamma) \geq (c - 2d)k$.*

Proposition 5.4. *Suppose that $4\epsilon < d \leq 1$ and that C is a cycle in the reduced graph Γ (including the case that C is a single edge). Then every cluster V_i contains a subcluster $V'_i \subseteq V_i$ of size $(1 - 2\epsilon)m$ such that $(V'_i, V'_j)_{G'}$ is $(2\epsilon, d - 3\epsilon)$-super-regular for every edge $V_iV_j \in E(C)$.*

We will use the following corollary of the well-known Blow-up lemma of Komlós, Sárközy, and Szemerédi [22]. An (ϵ, d, m)-*super-regular cycle* of order k is a graph obtained from taking a cycle C of order k (allowing for an edge to be a cycle of order 2) and replacing each vertex of C with a set of m vertices and replacing each edge of C with an (ϵ, d)-super-regular bipartite graph.

Corollary 5.5. *Let $k \geq 2$ and $0 < \frac{1}{m} \ll \epsilon, d$. An (ϵ, d, m)-super-regular cycle $\mathcal{C}$ of order k has the property that for all $u, v \in V(C)$, there exists a u, v-path of length at least $(m - 1)k$.*

Proof. Let $V_1, V_2, \ldots, V_k$ be a cyclic ordering (meaning $V_{1-1} = V_k$ and $V_{k+1} = V_1$) of the vertex set of $\mathcal{C}$ and let $u \in V_i$ and $v \in V_j$. Let $U_{i+1} = N(u) \cap V_{i+1}$ and $U_{j-1} = N(v) \cap V_{j-1}$ and note that $|U_{i+1}|, |U_{j-1}| \geq dm$. If each V_iV_{i+1} were complete, we could embed a path starting in U_{i+1} and ending in U_{j-1} which misses at most one vertex from each set. So we may use the blow-up lemma to embed a path $P \subseteq \mathcal{C}$ starting in U_{i+1} and ending in U_{j-1} which misses at most one vertex from each set. □

We use the following result of Gyárfás and Sárközy.

Theorem 5.6 (Gyárfás, Sárközy, [13])**.** *Let $\frac{1}{n_0} \ll \eta \leq 1/4$. If $\delta(G) \geq (3/4 + \eta)n$, then in every 2-coloring of G, there exists a monochromatic cycle of length at least $(2/3 - \eta)n$.*

We note that the statement of the above result in [13] is actually for paths instead of cycles; however, their proof implicitly gives the result for cycles. Furthermore, Benevides, Łuczak, Skokan, Scott, and White [2] proved a stronger version of Theorem 5.6 which says that if $\frac{1}{n_0} \ll \eta \leq 1/4$ and G is a graph on $n \geq n_0$ vertices with $\delta(G) \geq 3n/4$, then in every 2-coloring of G, there exists a monochromatic cycle of length at least $(2/3 - \eta)n$.

One of the key elements of Erdős and Galvin's proof that every 2-coloring of $K_{\mathbb{N}}$ contains a monochromatic path of strong upper density at least $1/(3+\sqrt{3})$ is the use of Theorem 1.1. One of the key elements of our proof of Theorem 1.7 is the following "robust" version of Theorem 1.1.

Corollary 5.7. *For all $0 < \eta \leq 1/4$ there exists $0 < \frac{1}{n_0} \ll \frac{1}{m} \ll \epsilon < 4d \ll \sigma$ such that every 2-colored graph K_n on $n \geq n_0$ vertices has a monochromatic (ϵ, d, m)-super-regular cycle C_1 on at least $(2/3 - \eta)n$ vertices.*

Proof. Apply Lemma 5.1 to the 2-colored K_n (with $\epsilon/2$ and $2d$) and then Proposition 5.3 to get 2-colored $(\epsilon/2, 2d)$-reduced graph Γ on k vertices with $\delta(\Gamma) \geq (1-4d)k \geq (3/4 + \epsilon/2)k$. Apply Theorem 5.6 to Γ to get a monochromatic cycle on at least $(2/3 - \epsilon/2)k$ vertices in Γ. Now use Proposition 5.4 to turn the cycle in Γ into a monochromatic (ϵ, d, m)-super-regular cycle in G on at least $(2/3 - \eta)n$ vertices. □

5.2 Specific Lemmas

Definition 5.8. *Let G be a graph on n vertices and let $\alpha \in [0, 1]$. We say that $X \subseteq G$ is an* α-connector *if, for all distinct $u, v \in V(X)$, there is a u, v-path $P \subseteq X$, such that $|V(P)| \geq \alpha n$.*

We say that that an α-connector X is maximal *if, for all $u \notin V(X)$, u has at most one neighbor in X.*

Lemma 5.9. *Let $0 < \frac{1}{n_0} \ll \epsilon \leq 2/3$. For all 2-colored complete graphs on $n \geq n_0$ vertices, there exists a monochromatic maximal α-connector X with $\alpha \geq 2/3 - \epsilon$.*

Proof. Apply Lemma 5.7 to get a monochromatic, say blue, super-regular cycle X_0 on at least $(2/3 - \epsilon/2)n$ vertices. Now for each i, let X_{i+1} be the set of vertices which have at least two blue edges into $\bigcup_{j=0}^{i} X_j$. Note that $X_i \subseteq X_{i+1}$. Now let $X = \bigcup_{i=0}^{\infty} X_i$.

If X is an α-connector, then by definition it will be maximal. So to see that X is an α-connector, let $u_i, v_j \in X$ where $u_i \in X_i$ and $v_j \in X_j$. By the definition of $X_1, X_2, \ldots$, it is possible to iteratively choose disjoint paths from u_i to some vertex $u_0 \in X_0$ and from v_j to some vertex $v_0 \in X_0$ with $u_0 \neq v_0$. Now using the fact that X_0 is a super-regular cycle on at least $(2/3 - \epsilon/2)n$ vertices, we find a blue path from u_0 to v_0 which uses nearly all of the vertices of X_0, i.e. a blue u_i, v_j-path on at least $(2/3 - \epsilon)n$ vertices. □

Lemma 5.10. *Let $0 < \frac{1}{n_0} \ll \epsilon \ll \alpha_1, \alpha_2 \leq 1$ where $\alpha_2 > 1/2 + \epsilon$ and let G be a 2-colored complete graph on a finite interval $V \subseteq \mathbb{N}$ with $|V| \geq n_0$. Let $V_1 \cup V_2$ be a partition of V such that V_1 is an initial segment of V, and both V_1 and V_2 have local density[4] at least $1 - \epsilon$. Suppose X_2 is a maximal blue α_2-connector in V_2.*

(i) *If $X_1 \subseteq V_1$ such that $|X_1|/|V_1| \geq \alpha_1$ and there is no blue matching of size 2 from X_1 to X_2, then for each $i \in [2]$, there is a red path P with both endpoints in X_i such that P has local density at least $(1 - \epsilon)^2 \alpha_1$.*

[4] See Section 2)

(ii) *If X_1 is a maximal red α_1-connector in V_1, then there exist $u_i, v_i \in X_i$ for some $i \in [2]$ and red and blue u_i, v_i-paths P_1 and P_2 respectively, such that P_j has local density at least $(1-\epsilon)^2\alpha_j$ for each $j \in [2]$.*

Proof. (i) Note that since V_2 has local density at least $1-\epsilon$, we have $|V_1| \leq \epsilon|V_2|$. Then, $|X_2| > |X_1|+|V_2 \setminus X_2|$, and it follows that there is an initial segment I of V_2 such that $|X_1|+|(V_2 \setminus X_2) \cap I| = |X_2 \cap I|$. Let $Y_1 = X_1 \cup ((V_2 \setminus X_2) \cap I)$ and $Y_2 = X_2 \cap I$. By the assumption and the fact that X_2 is a maximal α_2-connector in V_2, the bipartite graph $[Y_1, Y_2]$ has no blue matching of size 2 and thus by deleting at most one vertex from either Y_1 or Y_2, we have a bipartite graph which is complete in red. It follows that for each i we can find a red path P_i with both endpoints in Y_i which misses at most 3 vertices from $Y_1 \cup Y_2 = X_1 \cup I$. Then P_i has local density at least

$$\frac{|X_1|+|I|-3}{\min(V)+|V_1|+|I|} \geq \frac{|X_1|-3}{\min(V)+|V_1|} \geq (1-\epsilon)\frac{|X_1|-3}{|V_1|} \geq (1-\epsilon)^2\alpha_2.$$

(ii) Case 1: There is a blue matching $\{u_1u_2, v_1v_2\}$ of size 2 between X_1 and X_2. We construct P_1 and P_2 as follows. Since X_1 is a red α_1-connector inside $G[V_1]$, we may choose a red path P_1 with endpoints u_1 and v_1 such that $|P_1| \geq \alpha_1 |V_1|$. Similarly, we may choose a blue path $P_2^* \subseteq X_2$ with endpoints u_2 and v_2, such that $|P_2^*| \geq \alpha_2 |V_2|$. It follows that $d_\ell(P_1) \geq (1-\epsilon)\alpha_1$ and $d_\ell(P_2^*) \geq (1-\epsilon)\alpha_2$. Now we let $u = u_1$, $v = v_1$, and $P_2 = u_1u_2P_2^*v_2v_1$.

Case 2: There is no blue matching of size 2 between X_1 and X_2. Then we apply Lemma 5.10.(i) to get a red path P_1 with endpoints $u, v \in X_2$ such that P_1 has local density at least $(1-\epsilon)^2\alpha_1$. Since X_2 is a blue α_2-connector, we may also choose a blue path P_2 with endpoints u and v and $|P_2| \geq \alpha_2 |V_2|$, and it follows that the local density of P_2 is at least $(1-\epsilon)\alpha_2$. □

5.3 Main result

Proof of Theorem 1.7. Fix some sequence $\epsilon_n > 0$ tending to zero, and partition $\mathbb{N}$ into a sequence A_n of fast-growing intervals such that each $|A_n|$ satisfies the conclusion of Lemma 5.9 with $\epsilon = \epsilon_n$, and also such that the local density of A_n is at least $1-\epsilon_n$. By Lemma 5.9, for each n we may find a maximal α_n-connector X_n in A_n of some color χ_n, with $\alpha_n \geq 2/3 - \epsilon_n$. (See Figure 2 in each of the cases below)



(a) Case 1a (b) Case 1b

(c) Case 2

Figure 2: Cases in the proof of Theorem 1.7

Case 1: χ_n is eventually constant. Say, without loss of generality, that χ_n is eventually blue.

Case 1(a): There is an infinite sequence n_i in $\mathbb{N}$ such that for each i, there is a blue matching of size 2 between vertices in X_{n_i} and $X_{n_{i+1}}$. Using these matchings we may choose distinct vertices $u_i, v_i \in X_{n_i}$ such that for each i, the edge v_iu_{i+1} is blue. By Definition 5.8, we may choose blue

paths $P_i \subseteq X_{n_i}$, with endpoints u_i and v_i, such that $|P_i| \geq (2/3 - \epsilon_{n_i})\,|A_{n_i}|$, and hence the local density of P_i is at least $(2/3 - \epsilon_{n_i})(1 - \epsilon_{n_i})$. Then by joining the paths P_i with the edges $v_i u_{i+1}$, we get a blue path P with $\bar{d}_s(P) \geq 2/3$.

Case 1(b): There is an infinite sequence n_i in $\mathbb{N}$ such that for each i and for each $m > n_i$, there is no blue matching of size 2 between vertices in X_{n_i} and X_m. We may assume that for each i, $n_{i+1} > n_i + 1$. Note that the induced graph on $A_{n_i} \cup A_{n_i+1}$ satisfies the hypotheses of Lemma 5.10.(i). So there exists a red path P_i having both endpoints in A_{n_i} such that the local density of P_i is at least $(1 - \epsilon_{n_i})^3(2/3 - \epsilon_{n_i})$. Now by joining the P_i's with the edges $v_i u_{i+1}$, we get a red path P with $\bar{d}_s(P) \geq 2/3$.

Case 2: χ_n is not eventually constant. Then there is an infinite set $M \subseteq \mathbb{N}$ such that for all $n \in M$, $\chi_n \neq \chi_{n+1}$. By Lemma 5.10.(ii), for each $n \in M$ we may choose vertices $u_n, v_n \in X_n \cup X_{n+1}$ and monochromatic paths P_n and Q_n such that

(i) P_n and Q_n both have endpoints u_n, v_n,

(ii) P_n is blue and Q_n is red, and

(iii) each of P_n, Q_n has local density at least

$$(1 - \epsilon_n)^2(2/3 - \epsilon_n).$$

By Ramsey's theorem, we may find an infinite sequence n_i in M such that for each i, the edge $v_{n_i} u_{n_{i+1}}$ is the same color ρ. Using a combination of these edges and either the P_{n_i}'s or the Q_{n_i}'s, we may construct an infinite path P with color ρ, and

$$\bar{d}_s(P) \geq \limsup_{i \to \infty} (1 - \epsilon_{n_i})^2(2/3 - \epsilon_{n_i}) = 2/3,$$

which completes the proof □

6 Connected subgraphs

6.1 Covering/Partitioning by connected subgraphs

In this subsection we prove Theorem 6.1 which has as an immediate corollary a result about upper density of monochromatic connected subgraphs. While this corollary is superseded by the stronger result in the next subsection, we include it here both to show how a covering result can imply an upper density result and because Theorem 6.1 may be of independent interest.

Theorem 6.1. *For every 3-coloring of $K_{\mathbb{N}}$, either*

(i) *there exists a partition of $\mathbb{N}$ by at most two monochromatic connected subgraphs, or*

(ii) *every maximal[5] monochromatic connected subgraph is co-finite and there exists a cover of $\mathbb{N}$ by two monochromatic connected subgraphs T_1 and T_2 so that $V(T_1) \cap V(T_2)$ is finite.*

Proof. Let G be a maximal monochromatic connected subgraph and without loss of generality suppose it is green. If possible, choose G so that $\mathbb{N} \setminus V(G)$ is infinite, but otherwise choose G so that $\mathbb{N} \setminus V(G)$ is minimized. Let $U := \mathbb{N} \setminus V(G)$. If $|U| \leq 2$, then we have a partition of $\mathbb{N}$ into one green connected subgraph and the monochromatic subgraph induced by U; so suppose $|U| \geq 3$. Every edge from U to G is either red or blue, so choose a connected component R, say red, in the U, G-bipartite graph so that $V(R) \cap U$ is maximal and note that by the maximality of G, we

[5]Here maximal means not properly contained in a monochromatic connected subgraph of any color.

have $V(G) \setminus V(R) \neq \emptyset$. If $U \setminus V(R) \neq \emptyset$ then every edge from $V(G) \cap V(R)$ to $U \setminus V(R)$ is blue and every edge from $V(G) \setminus V(R)$ to $V(R) \cap U$ is blue, giving us a partition into two connected blue subgraphs. So suppose $U \setminus V(R) = \emptyset$. In this case every edge from U to $V(G) \setminus V(R)$ is blue giving us a blue connected subgraph B, and thus any two of $\{G, R, B\}$ forms a cover of $\mathbb{N}$.

Let

$$X' = \{v \in V(R) \cap V(G) : \text{ every edge from } v \text{ to } U \text{ is red}\} \text{ and } X = (V(R) \cap V(G)) \setminus X'.$$

Suppose that U is infinite, or U is finite and $|X| < |U|$. If $X' = \emptyset$, then $\mathbb{N} = V(B) \cup X$ induces a connected blue subgraph and we are done, so suppose not. By the assumption, we may choose an injective function f from X to U which is not surjective. Let $X_1 = \{v \in X : \{v, f(v)\} \text{ is blue}\}$ and let $X_2 = X \setminus X_1$ and without loss of generality suppose $X_1 \neq \emptyset$ (by defining X_1 as the red images if necessary). Let U_1 be the image of X_1 under f and let $U_2 = U \setminus U_1$ which is non-empty since f is not surjective. We now see that $X' \cup X_1 \cup U_1$ induces a blue connected subgraph and $X_2 \cup (V(B) \cap V(G)) \cup U_2$ induces a red connected subgraph, giving the desired partition.

We are left with the case that U is finite and $|X| \geq |U| \geq 3$. In this case, R and G form a cover of $\mathbb{N}$ in which $V(R) \cap V(G)$ is finite. □

Corollary 6.2. *For every 3-coloring of $K_{\mathbb{N}}$, there exists a monochromatic connected subgraph T such that $\bar{d}(T) \geq 1/2$.*

For finite graphs, Tuza [30] showed that for $r \in \{4, 5\}$, any r-colored complete graph can be covered with at most $r - 1$ monochromatic connected subgraphs. While it is likely that Tuza's argument extends to infinite graphs as in Theorem 6.1, we don't explore that here.

It is mentioned in [8] that Nagy and Szentmiklóssy proved a partition version of Theorem 6.1, but to our knowledge, the proof has never appeared in print. It would be interesting to recover this proof, and we note that our proof above would give a partition if the following was true.

Conjecture 6.3. *Let G be a complete bipartite graph with parts U, X, such that U is finite and $|X| \geq |U| \geq 3$ where X is possibly infinite. Suppose G is 2-colored in such a way that every vertex in X is incident with at least one edge of each color. Then there exists a collection of connected non-trivial subgraphs $B_1, \ldots, B_s$, and $R_1, \ldots, R_t$ with $s, t \geq 1$, such that each B_i is blue, each R_i is red, and $\bigcup_i \{B_i \cap X\} \cup \bigcup_j \{R_j \cap X\}$ is a partition of X.*

6.2 Strong upper density of connected subgraphs

First we give the short proof of the 2-color case.

Proof of Theorem 1.11. Partition $\mathbb{N}$ into fast-growing intervals A_n. By the remark of Erdős and Rado, in each interval A_i, there is a spanning monochromatic connected subgraph T_i. By the pigeonhole principle, there exists an infinite subsequence of say blue subgraphs $T_{n_1}, T_{n_2}, \ldots$. By Ramsey's theorem, either there exists a subsequence $T_{k_1}, T_{k_2}, \ldots$ such that for all i, there is a blue edge from T_{k_i} to $T_{k_{i+1}}$, in which case we have a blue connected subgraph with strong upper density 1. Otherwise, there exists a subsequence $T_{k_1}, T_{k_2}, \ldots$ such that for all i and all $j > i$, every edge from T_{k_i} to T_{k_j} is red. In this case we get a red connected subgraph of strong upper density 1 (actually, we can get a red star of strong upper density 1 with center in T_{k_1} and every other vertex in $T_{k_2} \cup T_{k_3} \cup \ldots$ as a leaf). □

In preparation for the 3-color case, we first prove a few lemmas.

Lemma 6.4. *In any 3-coloring of K_n,*

(i) *there exists a monochromatic connected subgraph on n vertices, or*

(ii) *there exists a partition $\{X, Y, Z, W\}$ of $[n]$ and monochromatic connected subgraphs $B_1 = X \cup Y$, $B_2 = Z \cup W$, $R_1 = X \cup Z$, $R_2 = Y \cup W$, $\{G_1, G_2\} = \{X \cup W, Y \cup Z\}$, and $|B_1|, |R_1|, |G_1| \geq n/2$, or*

(iii) *there exist monochromatic connected subgraphs, B, R, G such that any two of $\{B, R, G\}$ cover $[n]$ and $|B| \geq 2n/3$, $|R| \geq n - |R|/2 \geq n/2$, and $|G| \geq 2n - |B| - |R|$.*

Proof. Suppose B is the largest monochromatic, say blue, connected subgraph and set $U = [n] \setminus B$. Either $|B| = n$ in which case (i) is satisfied, or there is a monochromatic component R such that $|R \cap B| \geq |B|/2$ and $|R \cap U| \geq |U|/2$; note that by the maximality of B, we have $B \setminus R \neq \emptyset$.

(ii) If $U \setminus R \neq \emptyset$, then $[B \cap R, U \setminus R]$ and $[B \setminus R, U \cap R]$ are complete in green, $[B \setminus R, U \setminus R]$ is complete in red, and $[U \cap R, U \setminus R]$ and $[B \cap R, B \setminus R]$ are complete in blue. For each color there are two components of that color which partition the vertex set and thus there is a component of each color which has at least $n/2$ vertices.

(iii) If $U \setminus R = \emptyset$, then since $|B| \geq |R| \geq n - |B| + |B|/2 = n - |B|/2$, we have $|B| \geq 2n/3$ and the bipartite graph between $B \setminus R$ and U is complete in green, which means there is a green connected subgraph G with $|G| \geq n - |B| + n - |R| = 2n - |B| - |R|$. □

Corollary 6.5. *For each 3-coloring of K_n, we either have*

(i) *a monochromatic connected subgraph on n vertices, or*

(ii) *monochromatic connected components B, R and G in different colors, each with at least $n/2$ vertices, such that $B \cap R \cap G \neq \emptyset$, or*

(iii) *there are disjoint sets X, Y, Z such that $B = X \cup Y$, $R = Y \cup Z$, and $G = X \cup Z$ are contained in monochromatic components of different colors, at least two of B, R, G have at least $n/2$ vertices, and $|B \cup R \cup G| \geq 3n/4$.*

Proof. (i) corresponds to Lemma 6.4.(i).

(ii) corresponds to Lemma 6.4.(ii) in the case where $B_1 \cap R_1 \cap G_1 \neq \emptyset$.

(iii) corresponds to either Lemma 6.4.(iii), or Lemma 6.4.(ii) in the case where $B_1 \cap R_1 \cap G_1 = \emptyset$. If $B_1 \cap R_1 \cap G_1 = \emptyset$, then either $|B_1 \cap R_1| \leq n/4$ in which case $|B_1 \cup R_1| \geq 3n/4$, or $|B_1 \cap R_1| > n/4$ in which case $|G_1 \cup B_1|, |G_1 \cup R_1| > 3n/4$. □

Lemma 6.6. *Let G be a complete tripartite graph on n vertices. In every 2-coloring of G, there exists a monochromatic connected component with more than $n/2$ vertices.*

Proof. Let X_1, X_2, X_3 be the tripartition of the vertex set. Let B be a monochromatic connected component of maximal size, say with color blue. Then every edge between $X_i \setminus B$ and $X_j \cap B$ is red, for each pair $i \neq j$. It follows that B must contain at least two of X_1, X_2, X_3; otherwise there is a red connected component which strictly contains B. Say without loss of generality that $X_1, X_2 \subseteq B$. If $X_3 \subseteq B$ too, then we're done; otherwise there is a red component R containing $X_3 \setminus B$, X_1, and X_2. Hence whichever of B or R contains more of X_3 must contain greater than $n/2$ vertices. □

Proof of Theorem 1.12. Partition $\mathbb{N}$ into fast-growing intervals $I_1, I_2, \ldots$. By Corollary 6.5, the coloring in each interval I_n falls into cases (i), (ii) or (iii). By the pigeonhole principle there are infinitely many intervals of the same type. Let J_n enumerate them. Throughout the proof, we

will repeatedly have to pass to a further subsequence of the J_n's; to keep the notation simple, we will not change the name, and simply call each subsequence J_n.

Case 1: Each J_n is type (i), i.e. has a monochromatic connected component covering the entire interval. Without loss of generality, by passing to a further subsequence, we may assume that the color of this component is the same, say blue. If there is a infinite blue path (i.e. a sequence of blue edges $v_1u_2, v_2u_3, v_3u_4, \ldots$ with $u_i, v_i \in J_i$) through these intervals, then we are done, so after passing to a subsequence we may assume that all edges between two different J_n's are colored red and green. We consider J_n, J_{n+1}, and J_{n+2} and note that by Lemma 6.6 either there is a red connected subgraph or a green connected subgraph which has density (in $J_n \cup J_{n+1} \cup J_{n+2}$) at least $1/2$; suppose that, infinitely often, it is green. Now look at the edges just between these green components. We already know the edges between them are only red or green, so either there is an infinite green path in which case we get a green connected subgraph of strong upper density at least $1/2$, or we can pass to a subsequence which only consists of red edges in which case we get a red connected subgraph of strong upper density at least $1/2$.

Case 2: There are infinitely many intervals of type (ii). Let B_n, R_n and G_n be the monochromatic components of J_n as in Corollary 6.5. Fix vertices $v_n \in B_n \cap R_n \cap G_n$. Using Ramsey's theorem to pass to a subsequence, we may assume that each of the edges v_nv_{n+1} has the same color, say blue; then we have a blue connected subgraph with strong upper density at least $1/2$.

Case 3: There are infinitely many intervals of type (iii). Let X_n, Y_n, Z_n be the subsets of J_n as in this case; i.e. X_n, Y_n and Z_n are pairwise disjoint, the sets $B_n = X_n \cup Y_n$, $R_n = Y_n \cup Z_n$, and $G_n = X_n \cup Z_n$ are contained in monochromatic components of different colors, at least two of B_n, R_n, G_n have size at least $|J_n|/2$, and $|X_n \cup Y_n \cup Z_n| \geq 3|J_n|/4$. We may assume without loss of generality that B_n is connected in blue, R_n in red, and G_n in green; moreover, by passing to an infinite subsequence we may assume that $|B_n|, |R_n| \geq |J_n|/2$.

Suppose there is an infinite path of blue edges between B_n's; then we have a blue connected subgraph with strong upper density at least $1/2$. Then by passing to a subsequence, we may assume that there are no blue edges between any two B_n's. Similarly we may assume there are no red edges between any two R_n's; in particular, this implies that every edge between two Y_n's is green.

Now consider the edges between Y_m and X_n, for $m < n$. We know already that no such edges are blue. Suppose, on the other hand, that there is an infinite subsequence n_i such that there is a green edge between Y_{n_i} and $X_{n_{i+1}}$ for each i; then there is a green connected subgraph containing $X_{n_i} \cup Y_{n_i} \cup Z_{n_i}$ for each i, and hence we are done. So we may assume no such subsequence exists, and by passing to an infinite subsequence again this means that every edge from Y_m to X_n, for $m < n$, is red.

By similar logic, we either get a monochromatic connected subgraph with strong upper density at least $1/2$, or we may make the following deductions in the following order (including the two observations already made above), passing to an infinite subsequence at every stage:

(i) Every edge from Y_m to Y_n is green for every $m < n$.

(ii) Every edge from Y_m to X_n is red for every $m < n$.

(iii) Every edge from Y_m to Z_n is blue for every $m < n$.

(iv) Every edge from X_m to Y_n is green for every $m < n$.

(v) Every edge from Z_m to Y_n is green for every $m < n$.

(vi) Every edge from X_m to X_n is red for every $m < n$.

(vii) Every edge from Z_m to Z_n is blue for every $m < n$.

(viii) Every edge from X_m to Z_n is blue for every $m < n$.

(ix) Every edge from Z_m to X_n is red for every $m < n$.



Figure 3: A part of the infinite subsequence after making the deductions above. The red edges are solid, the blue edges are dashed, and the green edges are dotted.

To finish the proof, we must consider what happens between the intervals J_n. Let $J_n = [u_n, v_n]$. Note that each color has exactly two components in the graph restricted to $(X_1 \cup Y_1 \cup Z_1) \cup \cdots \cup (X_n \cup Y_n \cup Z_n)$, and these components together have density approaching $3/4$ as $n \to \infty$; e.g. $X_n \cup Y_n$ and Z_n are both contained in blue connected components.

For each $k \in (v_n, v_{n+1}]$, let B_k^* be the union of the two (not necessarily distinct) blue components containing $X_n \cup Y_n$ and Z_n, respectively, in the graph restricted to $\{1, \ldots, k\}$. Similarly we define R_k^* to be the union of the red components containing $Y_n \cup Z_n$ and X_n, and G_k^* the union of the green components containing $X_n \cup Z_n$ and Y_n. We call $k \in \mathbb{N}$ a *blue-connector* if B_k^* is connected in blue; similarly, we call k a red-connector or green-connector if R_k^* or G_k^* are connected in their respective colors. Note that there are connectors of each color in J_{n+1}. Our challenge is to show that there is a connector k in the interval $(v_n, v_{n+1}]$ whose corresponding connected subgraph has density at least $1/2$ in $\{1, \ldots, k\}$. Then, infinitely often, its color will be the same, and we have found the desired connected subgraph with strong upper density at least $1/2$.

Note that if k is not a connector of any color, then k must be in at least two of the sets B_k^*, R_k^*, G_k^*. Now let k_n be the first connector in the interval $(v_n, v_{n+1}]$. Since no vertex before k_n was a connector of any color, we have

$$\left|B_{k_n}^*\right| + \left|R_{k_n}^*\right| + \left|G_{k_n}^*\right| \geq 3\left|X_n \cup Y_n \cup Z_n\right| + 2(k_n - v_n) \geq \frac{9}{4}(v_n - u_n) + 2(k_n - v_n) \geq 2k_n.$$

(We have $9/4(v_n - u_n) \geq 2v_n$ as long as $v_n \geq 9u_n$, which we may fold into our assumption that $|J_n| \to \infty$ rapidly.) Say k_n is a blue connector. If $\left|B_{k_n}^*\right| \geq k_n/2$, then we are done. So suppose $\left|B_{k_n}^*\right| = k_n/2 - m$. Then $\left|R_{k_n}^*\right| + \left|G_{k_n}^*\right| \geq 3k_n/2 + m$, and thus $\min\{R_{k_n}^*, G_{k_n}^*\} \geq k_n/2 + m$. Now let ℓ_n be the next connector of a different color. If $\left|B_{\ell_n}^*\right| \geq \ell_n/2$, then again we are done; so suppose this is not true, i.e. fewer than $(\ell_n - k_n)/2 + m$ of the vertices in $(k_n, \ell_n]$ are in $B_{\ell_n}^*$. Then there are more than $(\ell_n - k_n)/2 - m$ vertices in (k_n, ℓ_n) which have no blue edges to $X_n \cup Y_n \cup Z_n$. On the other hand, since these vertices are not connectors in red or green, they must have edges of both colors back to $X_n \cup Y_n \cup Z_n$, and hence are included in both $R_{\ell_n}^*$ and $G_{\ell_n}^*$. Now say ℓ_n is a red connector; then we have

$$\left|R_{\ell_n}^*\right| \geq k_n/2 + m + (\ell_n - k_n)/2 - m = \ell_n/2,$$

which gives us the desired connector with sufficiently large density. □

7 0-dense graphs

Proposition 7.1. *Let $h : \mathbb{N} \to \mathbb{N}$, where $h(n) \to \infty$. There exists a 2-coloring of $K_{\mathbb{N}}$ such that if G is a graph in which the largest independent set has size $i < \infty$, then the vertices of any monochromatic copy of G must eventually grow faster than h. In particular, any graph with bounded independence number is 0-dense.*

Proof. Let $\mathbb{N} = A_1 \cup A_2 \cup \cdots \cup A_n \cup \cdots$ be a partition of $\mathbb{N}$ into consecutive intervals, such that $|A_n| = nh(n)$. Color the edges inside the intervals blue, and the edges between intervals red. Let G be an infinite graph with independence number $i < \infty$. Then if V is the vertex set of a monochromatic (necessarily red) copy of G, it follows that $|V \cap A_n| \leq i$ for each n, and from this that the vertices of V eventually grow faster than h. □

Let $\mathcal{T}^+$ be the transitive tournament on $\mathbb{N}$ such that for all $i, j \in \mathbb{N}$, $(i, j) \in \mathcal{T}^+$ if and only if $i < j$. Let $\mathcal{T}^-$ be the transitive tournament on $\mathbb{N}$ such that for all $i, j \in \mathbb{N}$, $(i, j) \in \mathcal{T}^-$ if and only if $i > j$.

Definition 7.2. *We define two classes of countably infinite directed graphs, $\mathscr{F}^+$ and $\mathscr{F}^-$, as follows. $\mathscr{F}^\pm$ (we use the symbol $\pm$ to define them simultaneously) is defined to be the class of countably infinite graphs G such that there is a partition $V(G) = A \cup B$ into infinite sets satisfying*

(i) *$A \cong \mathcal{T}^\pm$ and B is independent,*

(ii) *for each $a \in A$, $d^\mp(a, B)$ is finite,*

(iii) *for each $b \in B$, $d^\mp(b, A) = 0$.*

Theorem 7.3. *Let $\mathscr{U}$ be a nonprincipal ultrafilter. Let T be a tournament on $\mathbb{N}$, and let a k-coloring of T be given. Then the following is satisfied for at least one $s \in \{+, -\}$: for every $G \in \mathscr{F}^s$ there is an $A \in \mathscr{U}$ such that there is a monochromatic isomorphic copy of G in T with vertex set A.*

Proof. Fix T and a k-coloring of T. For each $v \in \mathbb{N}$ we have either $N_T^+(v) \in \mathscr{U}$ or $N_T^-(v) \in \mathscr{U}$; put $v \in S^+$ in the first case and $v \in S^-$ in the second case. Then the two sets $S^\pm$ partition $\mathbb{N}$ and hence one must be in $\mathscr{U}$. We will show that, if $S^\pm \in \mathscr{U}$, then for all $G \in \mathscr{F}^\pm$ there is a monochromatic copy of G in $\mathscr{U}$. We will do this in the case $S^+ \in \mathscr{U}$; the other case is symmetric.

So we assume $S^+ \in \mathscr{U}$. As above we partition S^+ into sets S_i, $i = 1, \ldots, k$, so that $v \in S_i$ if and only if $N_i^+(v) \in \mathscr{U}$. Since these sets partition S^+, one must be in $\mathscr{U}$; say without loss of generality that $S_1 \in \mathscr{U}$.

Fix $G \in \mathscr{F}^+$, and let $V(G) = A \cup B$ be a partition as in the definition of $\mathscr{F}^+$. Let a_i and b_i ($i \in \mathbb{N}$) enumerate the elements of A and B respectively, and choose a function $f : \mathbb{N} \to \mathbb{N}$ satisfying

$$N^-(a_i) \cap B \subseteq \{b_j \mid j \leq f(i)\}$$

(To avoid degenerate cases we also require $f(i) \to \infty$ as $i \to \infty$.)

Now we recursively choose vertices $x_1, x_2, \ldots$ and $y_1, y_2, \ldots$ from S_1 so that

(i) for all $i < j$, (x_i, x_j) has color 1,

(ii) for all i and $j \leq f(i)$, (y_j, x_i) has color 1, and

(iii) the sets $X = \{x_i \mid i \in \mathbb{N}\}$ and $Y = \{y_i \mid i \in \mathbb{N}\}$ partition S_1.

To begin the recursion, let $y_1, \ldots, y_{f(1)}$ be the first $f(1)$-many vertices of S_1. Let $A_i = N_1^+(y_i)$ for each $i \leq f(1)$; then each A_i is in $\mathscr{U}$, and hence there are infinitely-many x in $A_1 \cap \cdots \cap A_{f(1)} \cap S_1$; let x_1 be the minimal such x greater than $y_1, \ldots, y_{f(1)}$.

Now suppose we've chosen $x_1, \ldots, x_n$ and $y_1, \ldots, y_{f(n)}$. Let $y_{f(n)+1}, \ldots, y_{f(n+1)}$ be the next $(f(n+1) - f(n))$-many vertices of S_1 not so far listed. As in the base case, if we let $A_i = N_1^+(y_i)$ for each $i \leq f(n+1)$ and $B_j = N_1^+(x_j)$ for each $j \leq n$, then the set

$$S_1 \cap A_1 \cap \cdots \cap A_{f(n+1)} \cap B_1 \cap \cdots \cap B_n$$

is in $\mathscr{U}$, and hence we may choose x_{n+1} to be some element of this set larger than all x_i's and y_i's listed so far.

Given sets X and Y as described above, we can clearly find a monochromatic copy of G with vertex set S_1. This completes the proof. □

An ultrafilter $\mathscr{U}$ is called *Ramsey* if for every coloring of $K_{\mathbb{N}}$ in finitely-many colors, there is an $A \in \mathscr{U}$ such that the complete graph on A is monochromatic. (See [21] Chapter 17 for other properties of Ramsey ultrafilters.) Hence if $\mathscr{U}$ is Ramsey, then any (oriented) graph satisfies the conclusions of Theorem 7.3; however, it is known that the existence of Ramsey ultrafilters is independent of ZFC (see [31]). Moreover, every Ramsey ultrafilter necessarily contains sets whose elements grow arbitrarily fast; this is implied by the results of [5] but can also be proven directly.

On the other hand, it is well known that the subsets of $\mathbb{N}$ of density 1 form a filter and that any filter can be extended to an ultrafilter (by Zorn's lemma). So there exists an ultrafilter $\mathscr{U}$ such that every $A \in \mathscr{U}$ has positive upper density (see [21] Chapter 17, Problem 7). Applying Theorem 7.3 with this special ultrafilter $\mathscr{U}$, we get the following corollaries;

Corollary 7.4. *Let $P = (p_1, p_2, \ldots)$ be a directed path with infinitely many switches. Then for any r-coloring of a tournament on $\mathbb{N}$ there is a monochromatic copy of P with positive upper density.*

Say that a graph G is *half-locally finite* if G contains an infinite independent set B and that for all $v \in V(G) \setminus B$, $d(v, B)$ is finite. (Note that $G[V(G) \setminus B]$ could induce a complete graph for instance). Equivalently, G is half-locally finite if $G \subseteq F \in \mathscr{F}$, where $\mathscr{F}$ is the family of undirected graphs obtained by removing the orientations from say $\mathscr{F}^+$.

Corollary 7.5. *Let G be a half-locally finite graph. Then for any r-coloring of $K_{\mathbb{N}}$, there is a monochromatic copy of G with positive upper density.*

8 Open problems and concluding remarks

8.1 Paths

We conjecture the following extension of Theorem 1.6.

Conjecture 8.1. *Every 2-coloring of $K_{\mathbb{N}}$ contains a monochromatic path of upper density at least $8/9$. If true, this is best possible by Example 9.2.*

For finite graphs, Gyárfás, Ruszinkó, Sárközy, and Szemerédi [12] proved that for sufficiently large n, every 3-coloring of K_n contains a monochromatic path of length at least $n/2$. We conjecture the following analog for infinite graphs.

Conjecture 8.2. *Every 3-coloring of $K_{\mathbb{N}}$ contains a monochromatic path of upper density at least $1/2$. If true, this is best possible by Corollary 3.5.*

8.2 Two-way infinite paths, i.e. cycles

While we don't study this variant, it's likely that our result for strong upper density of paths extends to a two-way infinite version, which would be an analog of cycles for infinite graphs. However, since a result for two-way infinite paths does not imply a result for one-way infinite paths, we did not explore this possibility. On the other hand, for ordinary upper density, it's clear that our construction can be modified to give a two-way infinite path with upper density at least $3/4$.

8.3 Dense paths in both colors

We mention a variant of the problem which seems not to have been studied even for finite graphs. That is, it should be possible to extend Theorem 1.1 to say that in every 2-coloring of K_n, there exists a red path P_R with more than $\alpha n \geq 2n/3$ vertices and a blue path P_B with more than $2(1-\alpha)n$ vertices (even stronger would be the additional conclusion that $V(P_R) \cup V(P_B) = V(K_n)$). Again, it seems possible that such a result would extend to upper density and strong upper density versions, but we don't explore that possibility here.

8.4 Directed paths

For consistently oriented paths (equivalently paths with finitely many switches), the only case we weren't able to settle is the following.

Problem 8.3. *Does there exist a 2-coloring of $\vec{K}_{\mathbb{N}}$ in which every consistently oriented monochromatic path has upper density 0?*

We showed that if a path has infinitely many switches it is positively-dense, but is there an absolute lower bound? If we quantify how the switches are distributed, can we say something more precise?

Problem 8.4. *Is it true that for every $\epsilon > 0$, there exists a path P with infinitely many switches and a 2-coloring of $\vec{K}_n$ such that every monochromatic copy of P has upper density less than ϵ?*

Let f be an increasing function from $\mathbb{N}$ to $\mathbb{N}$ and let $P_f = v_1v_2v_3\ldots$ be a directed path such that $v_{f(i)}$ is a switch. Determine a lower bound on the upper density of monochromatic copy of P in $\vec{K}_n$. For instance if $f(i) = i$ for all $i \in \mathbb{N}$, then P_f is an anti-directed path and Rado's result implies that every 2-coloring of $\vec{K}_{\mathbb{N}}$ contains a monochromatic anti-directed path of upper density at least $1/2$.

8.5 Connected subgraphs

We conjecture the following infinite analog of Theorem 1.9, for which we provided the proof when $r \in \{2, 3\}$.

Conjecture 8.5. *For $r \geq 4$, every r-coloring of $K_{\mathbb{N}}$ contains a monochromatic connected subgraph T with $\bar{d}(T) \geq \bar{d}_S(T) \geq 1/(r-1)$. If true, this is best possible when $r-1$ is a prime power by Example 3.4.*

8.6 Relative density

We showed that consistently oriented paths are $(0,3)$-dense and graphs with bounded independence number are 0-dense, and in both cases we showed that no bound can be put on the growth of the vertices in a monochromatic copy. Is there a 0-dense graph for which some bound can be placed on the growth?

Problem 8.6. *Does there exist a graph G and a 2-coloring of $K_{\mathbb{N}}$ such that for every monochromatic copy of G, we have $\bar{d}(G) = 0$, but there exists an increasing unbounded function h such that there exists a monochromatic copy of G with $\bar{d}_h(G) = \limsup_{n\to\infty} \frac{h(|V(G)\cap[n]|)}{h(n)} > 0$?*

8.7 0-dense graphs

Chvátal, Rödl, Szemerédi, and Trotter [3] proved that for all $\Delta \in \mathbb{N}$, there exists $0 < c \leq 1$ such that if G is a graph on at most cn vertices with maximum degree at most Δ, then in every 2-coloring of K_n, there exists a monochromatic copy of G. We conjecture the following infinite analog of their result.

Conjecture 8.7. *For all $\Delta \in \mathbb{N}$ there exists $c > 0$ such that for every infinite graph G with maximum degree at most Δ and every 2-coloring of $K_{\mathbb{N}}$, there exists a monochromatic copy of G with $\bar{d}(G) \geq c$.*

We showed that if a graph G is half-locally finite, then it is positively-dense, but is there an absolute lower bound? If so, this would strongly imply the previous conjecture.

Problem 8.8. *For all $\epsilon > 0$, does there exists a half-locally finite graph G and a 2-coloring of $K_{\mathbb{N}}$ such that every monochromatic copy of G has upper density less than ϵ? If so, what about a locally finite graph G?*

We show that graphs with bounded independence number are 0-dense and half-locally finite graphs are positively-dense. It would be nice to have a complete characterization.

Problem 8.9. *Completely characterize which graphs G are positively-dense.*

8.8 Lower density

Problem 8.10. *Let G be a graph. Is it true that if there exists a 2-coloring of $K_{\mathbb{N}}$ such that for every monochromatic copy of G we have $\bar{d}(G) < 1$, then there exists 2-coloring of $K_{\mathbb{N}}$ such that for every monochromatic copy of G we have $\underline{d}(G) = 0$?*

8.9 Density results from partitioning results

In a recent paper, Elekes, D. Soukup, L. Soukup, and Szentmiklóssy [4], proved a variety of results about partitioning infinite (hyper)graphs. A number of these results have consequences for the density Ramsey problem studied in this paper. They prove the following three results:

(i) In any r-coloring of $K^k_{\mathbb{N}}$ (the complete k-uniform hypergraph on $\mathbb{N}$), there exists a partition of $\mathbb{N}$ into at most r monochromatic tight paths, and consequently there exists a monochromatic tight path with upper density at least $1/r$. The analogous problem for finite hypergraphs was studied in [14], [19] for loose cycles, and [18] for tight cycles.

(ii) In any 2-coloring of $K_{\mathbb{N}}$, there exists a partition of $\mathbb{N}$ into at most four monochromatic squares of paths and thus there exists a monochromatic square path of upper density at least

1/4. More generally, they showed that in any r-coloring of $K_{\mathbb{N}}$, there exists a partition of $\mathbb{N}$ into at most $r^{(k-1)r+1}$ monochromatic kth powers of paths and a finite set, and thus there exists a monochromatic kth power of a path with upper density at least $\frac{1}{r^{(k-1)r+1}}$. The analogous problem for squares of paths in 2-colored finite complete graphs was considered in [1].

(iii) In any r-coloring of the infinite random graph on $\mathbb{N}$ there exists a partition of $\mathbb{N}$ into at most r monochromatic paths and thus some monochromatic path has upper density at least $1/r$. The analogous problem for finite random graphs was considered in [24].

It would be interesting to improve any of these density results (as Erdős and Galvin improved the corollary of Rado's result) and to consider the corresponding strong upper density versions.

References

[1] P. Allen, B. Roberts, and J. Skokan. Ramsey numbers of squares of paths. *Electronic Notes in Discrete Mathematics* **49** (2015), 637–641.

[2] F. Benevides, T. Łuczak, J. Skokan, A. Scott, and M. White. Monochromatic cycles in 2-coloured graphs, *Combinatorics, Probability, and Computing*, **21** no. 1-2, (2012), 57–87.

[3] C. Chvatál, V. Rödl, E. Szemerédi, and W. T. Trotter. The Ramsey number of a graph with bounded maximum degree. *Journal of Combinatorial Theory, Series B* **34**, no. 3 (1983), 239–243.

[4] M. Elekes, D. Soukup, L. Soukup, and Z. Szentmiklóssy. Decompositions of edge-colored infinite complete graphs into monochromatic paths. arXiv:1502.04955, (2015).

[5] P. Erdős. Some remarks on Ramsay's theorem. Canadian Mathematical Bulletin, **7**, (1964), 619–622.

[6] P. Erdős and F. Galvin. Some Ramsey-type theorems. *Discrete mathematics* **87**, no. 3 (1991), 261–269.

[7] P. Erdős and F. Galvin. Monochromatic infinite paths. *Discrete Mathematics* **113**, no. 1 (1993), 59–70.

[8] P. Erdős, A. Gyárfás, and L. Pyber. Vertex coverings by monochromatic cycles and trees, *Journal of Combinatorial Theory Series B*, **51** (1991), 90–95.

[9] L. Gerencsér and A. Gyárfás. On Ramsey-type problems, *Ann. Sci. Budapest. Eötvös Sect. Math*, **10** (1967), 167–170.

[10] A. Gyárfás. Partition covers and blocking sets in hypergraphs. MTA SZTAKI Tanulmányok, **71**, 1977.

[11] A. Gyárfás and J. Lehel. A Ramsey-type problem in directed and bipartite graphs. *Periodica Mathematica Hungarica* **3**, no. 3-4 (1973), 299–304.

[12] A. Gyárfás, M. Ruszinkó, G. Sárközy, and E. Szemerédi. Three-color Ramsey numbers for paths. Combinatorica, **27**, no. 1 (2007), 35–69.

[13] A. Gyárfás and G. Sárközy. Star Versus Two Stripes Ramsey Numbers and a Conjecture of Schelp. *Combinatorics, Probability and Computing* **21**, no. 1-2 (2012), 179–186.

[14] A. Gyárfás, G. Sárközy, and E. Szemerédi. The Ramsey number of diamond-matchings and loose cycles in hypergraphs. *Electron. J. Combin* **15**, no. 1 (2008), # R126.

[15] I. Farah. Semiselective coideals. *Mathematika* **45**, no. 1 (1998), 79–103.

[16] M. Hrušàk. Combinatorics of filters and ideals. *Set theory and its applications* **533**, Contemp. Math., Amer. Math. Soc., Providence, RI (2011), 29–69.

[17] A.R.D Mathias. Happy families. *Ann. Math. Logic* **12**, no. 1 (1977), 59–111.

[18] P. Haxell, T. Łuczak, Y. Peng, V. Rödl, A. Ruciński, and J. Skokan. The Ramsey number for 3-uniform tight hypergraph cycles. *Combinatorics, Probability and Computing* **18**, no. 1-2 (2009), 165–203.

[19] P. Haxell, T. Łuczak, Y. Peng, V. Rödl, A. Ruciński, M. Simonovits and J. Skokan. The Ramsey number for hypergraph cycles I. *Journal of Combinatorial Theory, Series A* **113**, no. 1 (2006), 67–83.

[20] D. Kühn and D. Osthus. Embedding large subgraphs into dense graphs, arXiv:0901.3541, (2009).

[21] P. Komjáth and V. Totik. *Problems and theorems in classical set theory* Springer Science & Business Media (2006).

[22] J. Komlós, G. Sárközy, and E. Szemerédi. Blow-up lemma, *Combinatorica* **17**, no. 1 (1997), 109–123.

[23] J. Komlós and M. Simonovits. Szemerédi's regularity lemma and its applications in graph theory. *Bolyai Society Mathematical Studies 2, Combinatorics, Paul Erdős is Eighty (Vol. 2)*, Budapest (1996), 295–352.

[24] S. Letzter. Path Ramsey number for random graphs. *Combinatorics, Probability and Computing* **25**, no. 4 (2016), 612–622.

[25] A. Pokrovskiy. Partitioning edge-coloured complete graphs into monochromatic cycles and paths, *Journal of Combinatorial Theory, Series B* **106** (2014), 70–97.

[26] R. Rado. Monochromatic paths in graphs. *Ann. Discrete Math* **3** (1978), 191–194.

[27] F.P. Ramsey. On a problem of formal logic, *Proc. London Math. Soc., 2nd Ser.*, **30** (1930),264–286.

[28] H. Raynaud. Sur le circuit hamiltonien bi-coloré dans les graphes orientés. *Periodica Mathematica Hungarica* **3**, no. 3-4 (1973), 289–297.

[29] E. Szemerédi. Regular Partitions of Graphs. *Colloques Internationaux C.N.R.S - Problèmes Combinatoires et Théorie des Graphes*, **260** (1976), 399–401.

[30] Z. Tuza. Ryser's conjecture on transversals of r-partite hypergraphs. *Ars Combinatoria* **16** (1983), 201–209.

[31] E.L. Wimmers, The Shelah P-point independence theorem. *Israel Journal of Mathematics* **43**, no. 1 (1982), 28–48.

9 Appendix: Erdős-Galvin examples

The following two examples appear as Corollary 1.6 and Corollary 1.9 in [7]. As they are stated in more general terms in their paper, we reproduce them here in less general terms for the convenience of the reader.

Example 9.1. *There exists a 2-coloring of $K_{\mathbb{N}}$ such that for all monochromatic paths P, $\bar{d}_S(P) \leq 2/3$.*

Proof. Let A be the set of $n \in \mathbb{N}$ divisible by 3, and B its complement. Color every edge from A to B red; and color every edge from A to A, and from B to B, blue. There is clearly a blue path which covers all of B in order, and this path has strong upper density, and upper density, equal to $2/3$. Now suppose $P = (p_0, p_1, \ldots, p_n, \ldots)$ is any red path; say, without loss of generality, that $p_0 \in A$. Then for every n, $p_{2n} \in A$ and $p_{2n} \geq 3n$. Hence the density of $p_0, \ldots, p_{2n}$ inside any interval that contains this initial segment must be at most $2/3$. □

Example 9.2. *There exists a 2-coloring of $K_{\mathbb{N}}$ such that for all monochromatic paths P, $\bar{d}(P) \leq 8/9$.*

Proof. Let A_0, A_1, A_2 be a sequence of consecutive intervals in $\mathbb{N}$, with $|A_n| = 2^n$ for all $n \geq 0$. If n is even and $m \geq n$, color all edges between A_n and A_m red. If n is odd and $m \geq n$, color all edges between A_n and A_m blue (See Figure 4).



Figure 4: An illustration of the example defined above.

By symmetry, we can consider the upper density of a red path P, which we can see is maximized by alternating between vertices in A_0 and A_1, using the earliest vertices in A_1 possible, until A_0 has filled up; then move to A_2 and continue. So the upper density of P is

$$\lim_{n\to\infty} \frac{2 \cdot 2^0 + 2 \cdot 2^2 + \cdots + 2 \cdot 2^{2n}}{2^0 + 2^1 + \cdots + 2^{2n} + 2^{2n}} = \lim_{n\to\infty} \frac{2\sum_{i=0}^{n} \frac{1}{4^i}}{1 + \sum_{i=0}^{n} \frac{1}{2^i}} = \frac{2 \cdot \frac{4}{3}}{1+2} = 8/9.$$

□